

HEARTGARD® PLUS CHEWABLES

Rx

Merial

Parasiticide-Oral
(ivermectin/pyrantel)
NADA No.: 140-971

Active Ingredient(s): Each chewable tablet contains:

Color Coding	Ivermectin Content	Pyrantel Content
Blue	68 mcg	57 mg
Green	136 mcg	114 mg
Brown	272 mcg	227 mg

Indications: For use in dogs to prevent canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for a month (30 days) after infection and for the treatment and control of ascarids (*Toxocara canis*, *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*, *Ancylostoma braziliense*).

Dosage and Administration: Dosage: HEARTGARD® Plus (ivermectin/pyrantel) Chewables should be administered orally at monthly intervals at the recommended minimum dose level of 6 mcg of ivermectin per kilogram (2.72 mcg/lb) and 5 mg of pyrantel (as pamoate salt) per kg (2.27 mg/lb) of body weight. The recommended dosing schedule for prevention of canine heartworm disease and for the treatment and control of ascarids and hookworms is as follows:

Dog Weight	Chewables Per Month	Ivermectin Content	Pyrantel Content	Color Coding on Foil-Backing and Carton
Up to 25	1	68 mcg	57 mg	Blue
26 - 50	1	136 mcg	114 mg	Green
51 - 100	1	272 mcg	227 mg	Brown

HEARTGARD® Plus is recommended for dogs 6 weeks of age and older.

For dogs over 100 lb use the appropriate combination of these chewables.

Administration: Remove only one chewable at a time from the foil-backed blister card. Return the card with the remaining chewables to its box to protect the product from light. Because most dogs find HEARTGARD® Plus palatable, the product can be offered to the dog by hand. Alternatively, it may be added intact to a small amount of dog food. The chewable should be administered in a manner that encourages the dog to chew, rather than to swallow without chewing. Chewables may be broken into pieces and fed to dogs that normally swallow treats whole. Care should be taken that the dog consumes the complete dose, and treated animals should be observed for a few minutes after administration to ensure that part of the dose is not lost or rejected. If it is suspected that any of the dose has been lost, redosing is recommended.

HEARTGARD® Plus should be given at monthly intervals during the period of the year when mosquitoes (vectors), potentially carrying infective heartworm larvae, are active. The initial dose must be given within a month (30 days) after the dog's first exposure to mosquitoes. The final dose must be given within a month (30 days) after the dog's last exposure to mosquitoes.

When replacing another heartworm preventive product in a heartworm disease prevention program, the first dose of HEARTGARD® Plus must be given within a month (30 days) of the last dose of the former medication.

If the interval between doses exceeds a month (30 days), the efficacy of ivermectin can be reduced. Therefore, for optimal performance, the chewable must be given once a month on or about the same day of the month. If treatment is delayed, whether by a few days or many, immediate treatment with HEARTGARD® Plus and resumption of the recommended dosing regimen will minimize the opportunity for the development of adult heartworms.

Monthly treatment with HEARTGARD® Plus also provides effective treatment and control of ascarids (*T. canis*, *T. leonina*) and hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*). Clients should be advised of measures to be taken to prevent reinfection with intestinal parasites.

Efficacy: HEARTGARD® Plus Chewables, given orally using the recommended dose and regimen, are effective against the tissue larval stage of *D. immitis* for a month (30 days) after infection and, as a result, prevent the development of the adult stage. HEARTGARD® Plus Chewables are also effective against canine ascarids (*T.*

canis, *T. leonina*) and hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*).

Precaution(s): Store at controlled room temperature of 59°-86°F (15°-30°C). Protect product from light.

Caution(s): Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

All dogs should be tested for existing heartworm infection before starting treatment with HEARTGARD® Plus which is not effective against adult *D. immitis*. Infected dogs must be treated to remove adult heartworms and microfilariae before initiating a program with HEARTGARD® Plus.

While some microfilariae may be killed by the ivermectin in HEARTGARD® Plus at the recommended dose level, HEARTGARD® Plus is not effective for microfilariae clearance. A mild hypersensitivity-type reaction, presumably due to dead or dying microfilariae and particularly involving a transient diarrhea, has been observed in clinical trials with ivermectin alone after treatment of some dogs that have circulating microfilariae.

Warning(s): Keep this and all drugs out of the reach of children.

In case of ingestion by humans, clients should be advised to contact a physician immediately. Physicians may contact a Poison Control Center for advice concerning cases of ingestion by humans.

Adverse Reactions: In clinical field trials with HEARTGARD® Plus, vomiting or diarrhea within 24 hours of dosing was rarely observed (1.1% of administered doses). The following adverse reactions have been reported following the use of Heartgard®: Depression/lethargy, vomiting, anorexia, diarrhea, mydriasis, ataxia, staggering, convulsions and hypersalivation.

Trial Data: Safety: HEARTGARD® Plus has been shown to be bioequivalent to Heartgard®, with respect to the bioavailability of ivermectin. The dose regimens of HEARTGARD® Plus and Heartgard® are the same with regard to ivermectin (6 mcg/kg). Studies with ivermectin indicate that certain dogs of the Collie breed are more sensitive to the effects of ivermectin administered at elevated dose levels (more than 16 times the target use level) than dogs of other breeds. At elevated doses, sensitive dogs showed adverse reactions which included mydriasis, depression, ataxia, tremors, drooling, paresis, recumbency, excitability, stupor, coma and death. Heartgard® demonstrated no signs of toxicity at 10 times the recommended dose (60 mcg/kg) in sensitive Collies. Results of these trials and bioequivalency studies, support the safety of Heartgard® products in dogs, including Collies, when used as recommended.

HEARTGARD® Plus has shown a wide margin of safety at the recommended dose level in dogs, including pregnant or breeding bitches, stud dogs and puppies aged 6 or more weeks. In clinical trials, many commonly used flea collars, dips, shampoos, anthelmintics, antibiotics, vaccines and steroid preparations have been administered with HEARTGARD® Plus in a heartworm disease prevention program.

In one trial, where some pups had parvovirus, there was a marginal reduction in efficacy against intestinal nematodes, possibly due to a change in intestinal transit time.

Acceptability: In acceptability and field trials, HEARTGARD® Plus Chewables were shown to be an acceptable oral dosage form that was consumed at first offering by the majority of dogs.

Presentation: HEARTGARD® Plus is available in three dosage strengths (see Dosage section) for dogs of different weights. Each strength comes in convenient packages of 6, 9 and 12 chewables.

®HEARTGARD is a registered trademark of Merial.

7/98

Disclaimer: Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the product label or package insert. Compendium Code No.: 08491027